

Notes on the type of *Bubo virginianus* *sclariventris*

by Robert W. Dickerman

Received 8 October 2001; revision received 29 January 2004

In attempting to make order out of the variation he noted among the 50 pale Great Horned Owls *Bubo virginianus* from Ontario in the Royal Ontario Museum (ROM), Snyder (1961) divided them into more heavily and less heavily barred groups. Considering the less heavily marked birds to be *subarcticus*, he named the darker birds *B. v. sclariventris*, using a female taken on 26 February 1948 as the holotype. The condition of its gonads was not given. He apparently used only specimens in the ROM, and did not mention the existing types of *subarcticus* and *occidentalis*. The range of *sclariventris* was mapped by Karalus & Eckert (1974), Johnsgard (1988) and McGillivray (1989), and the name was used by James (1991) and Pittaway (1993).

Godfrey (1986) and Browning (1990) were unable to express an opinion on the validity of *sclariventris*, and it has not been included in any taxonomic work to date, including Houston *et al.* (1998) and König *et al.* (1999). Dickerman (1992) studied the series of 60 nesting-season specimens of the subarctic populations in the then National Museum of Canada, Ottawa, and found no distinctive subpopulations, but the Ottawa collection is weak in material from Ontario.

To help evaluate the validity of the name *sclariventris*, James A. Dick and S. Ross James of the ROM selected for me paratype 34,2,2,1 as being most similar to the holotype. I compared it with the types of *subarcticus* and *occidentalis* in the Academy of Natural Sciences in Philadelphia. Those two types differ, that of *occidentalis* being more heavily barred. Giving some allowance for possible spoiling of the specimens due to their different ages (1948 vs 1880), the paratype of *sclariventris* was a very close match to the holotype of *occidentalis*. Thus *B. v. sclariventris* Snyder is a junior synonym of *B. v. occidentalis* Stone, earlier considered to be a junior synonym of *B. v. subarcticus* Hoy (Stone 1897, Dickerman 1991).

In the absence of adequate nesting-season material, especially from the northern fringes of the range, *subarcticus* must be used for specimens showing a spectrum of variation in the extent of dark markings, and perhaps in the degree of buffy coloration (Dickerman 1992). If in the future *subarcticus* is subdivided, close attention must be paid to the differences between the types of *subarcticus* and *occidentalis*, both migrants, from Wisconsin and Iowa respectively.

Acknowledgments

I wish to especially thank J. A. Dick and S. R. James of the ROM for selecting the relevant paratype and for the loan thereof. My gratitude goes to the curators of the several museums where specimens were examined: Academy of Natural Sciences, Philadelphia; American Museum of Natural History, New York; and the National Museum of Natural History, Washington, D.C.

References:

- Browning, M. R. 1990 Taxa of two North American birds described from 1957–1987. *Proc. Biol. Soc. Wash.* 103: 432–451.
- Dickerman, R. W. 1991. On the validity of *Bubo virginianus occidentalis* Stone. *Auk* 108: 964–965.
- Dickerman, R. W. 1992. Additional specimens of the subarctic Great Horned Owl from New York. *Kingbird* 42: 73–75.
- Godfrey, W. E. 1986. *The Birds of Canada*. Revised edition. National Museum of Natural Sciences, (Canada), Ottawa.
- Houston, C. S., Smith, D. G. & Rohner, C. 1998. Great Horned Owl (*Bubo virginianus*). In Poole, A. & Gill, F. (eds.) *The birds of North America*, no. 372. The Birds of North America, Inc., Philadelphia.
- James, R. D. 1991. *Annotated checklist of the birds of Ontario*. Second edn. Life Sciences Miscellaneous Publications, Royal Ontario Museum, Toronto.
- Johnsgard, P. A. 1988. *North American owls, biology and natural history*. Smithsonian Institution Press, Washington, D.C.
- Karalus, K. E. & Eckert, A. W. 1974. *The owls of North America (north of Mexico), all of the species and subspecies illustrated in color and fully described*. Doubleday & Co., New York.
- König, C., Weick, F. & Becking, J.-H. 1999. *Owls, a guide to the owls of the world*. Pica Press, Robertsbridge.
- McGillivray, W. B. 1989. Geographic variation in size and reverse size dimorphism of the Great Horned Owl in North America. *Condor* 91: 777–786
- Pittaway, R. 1993. Recognizable forms, subspecies of the Great Horned Owl. *Ontario Birds* 11: 64–69.
- Stone, W. 1897. Proper name for the western Horned Owl of North America. *Amer. Nat.* 31: 236
- Snyder, L. L. 1961. On an unnamed population of the Great Horned Owl. *Contribution* 54: 1–7. Life Sciences Division, Royal Ontario Museum, Toronto

Address: R. W. Dickerman, Museum of Southwestern Biology, University of New Mexico, Albuquerque, New Mexico, 87131, U.S.A. Email: bobdickm@unm.edu

© British Ornithologists' Club 2004

Interesting distributional records and notes on the biology of bird species from a cloud forest reserve in north-west Ecuador

by Juan F. Freile & Jaime A. Chaves

Received 1 January 2003

Even in well-studied countries, such as Ecuador, new distributional records and range extensions are regularly reported as unexplored or poorly known areas are surveyed (e.g. Krabbe 1992, Krabbe *et al.* 1997, Freile 2001a). In this note we present new records, as well as latitudinal and altitudinal range extensions for several species, taking Ridgely & Greenfield (2001) as our baseline. Our records are based on studies carried out at Bosque Integral Otonga (hereafter Otonga), from March 1999 to February 2002 (see also Chaves 2001). Elevation is given within each species account. Taxonomy follows Ridgely & Greenfield (2001).